

## Review

## Neonatal Transitions in Social Behavior and Their Implications for Autism

Sarah Shultz,<sup>1,2,\*</sup> Ami Klin,<sup>1,2,3</sup> and Warren Jones<sup>1,2,3,\*</sup>

**Within the context of early infant–caregiver interaction, we review a series of pivotal transitions that occur within the first 6 months of typical infancy, with emphasis on behavior and brain mechanisms involved in preferential orientation towards, and interaction with, other people. Our goal in reviewing these transitions is to better understand how they may lay a necessary and/or sufficient groundwork for subsequent phases of development, and also to understand how the breakdown thereof, when development is atypical and those transitions become derailed, may instead yield disability. We review these developmental processes in light of recent studies documenting disruptions to early-emerging brain and behavior mechanisms in infants later diagnosed with autism spectrum disorder, shedding light on the brain–behavior pathogenesis of autism.**

**Neonatal Development in Context**

At birth, the behavioral repertoire of human newborns is limited [1,2]. Within just days, weeks, and months, however, early reflexes and reflex-like predispositions give way to complex volitional behavior: infants acquire new vocal abilities [3]; change their feeding, sleeping, and waking patterns [4–6]; and acquire new control of their eyes, neck, hands, and feet [7–12]. Brain size at birth, approximately one-third that of an adult's [13], doubles during the first year [14] and increases by another 35% by year 3 [15]. Synaptic density quadruples in year 1 and will be 150–200% greater than that of an adult by year 3 [16,17] (to then be pruned or selectively strengthened in iterations that continue throughout the lifespan [17,18]).

This enormous postnatal change, however, does not happen in a vacuum [19]; in the absence of near-constant care, the infant's survival would be nonviable [20,21]. The infant enters the world in a state of utter fragility, and requires a parent or primary caregiver to provide for survival needs. In playing that role, the caregiver serves as both partner and facilitator, matching his or her own behavior (in facial affect [22,23], vocal tone [24,25], and physical touch [26,27]) to the needs of the infant in a manner that serves as the foundation for further acquisition of abilities [27–29]. The behaviors of infant and caregiver are thus mutually adapted and mutually reinforcing [30]: the infant–caregiver dyad, iteratively changing itself through interaction, is an integral and inseparable unit of early development [31].

Within that context, we review a series of pivotal transitions in the first 6 months of typical infancy that depend upon the mutually reinforcing context of the infant–caregiver dyad, with emphasis on behavior and brain mechanisms thought to be involved in preferential orientation towards, and interaction with, other people. This topic is especially timely given recent reports demonstrating disruptions to early-emerging mechanisms of brain and behavior in infants later diagnosed with autism spectrum disorder (ASD) [32–34], highlighting early periods of infancy as key targets for future research. In reviewing this literature, we endeavored to link recent work to

**Highlights**

From the first moments of life, neonates exhibit a range of socially adaptive preferences and reflex-like responses that serve to orient their attention towards caregivers, as well as behaviors that serve as important signals to those caregivers.

Within the first 6 months of typical infancy, a series of pivotal transitions occur within the context of early infant–caregiver interaction, as initially spontaneous reflex-like responses transition into remarkably sensitive and contingent social action.

Recent reports suggest that these developmental transitions may be disrupted in autism spectrum disorder, opening a critical theoretical insight into understanding the brain–behavior pathogenesis of autism.

<sup>1</sup>Marcus Autism Center, Children's Healthcare of Atlanta, Atlanta, GA 30329, USA

<sup>2</sup>Division of Autism and Related Disabilities, Department of Pediatrics, Emory University School of Medicine, Atlanta, GA 30022, USA

<sup>3</sup>Center for Translational Social Neuroscience, Emory University, Atlanta, GA 30022, USA

\*Correspondence: sarah.shultz@emory.edu (S. Shultz) and warren.jones@emory.edu (W. Jones).

a broader set of seminal papers in neonatal behavior and development ([additional references can be found in online supplemental materials](#)). Only by intensively studying these periods and more fully understanding the growth trajectories that define normative development, will we be able to gain new mechanistic insights into the pathogenesis of autism and related disabilities.

### Mechanisms of Social Adaptive Action Present Shortly After Birth

The prolonged period of caregiver dependency and biological immaturity of human neonates, lasting years longer than that of other primate species [35], has often been noted as a critical aspect of human adaptive success [36]. More than simply providing a longer period of postnatal plasticity in which to refine brain and behavior [37], the initial fragility and prolonged maturation of the human newborn provide an ineluctable 'initial task' (that of survival) as well as a highly specific context in which that task will be solved. That context, the specific setting in which changes in newborn brain and behavior will take place, is defined by nearly continuous interaction with other people.


#### Preferential Orientation and Reflexive Engagement

Shortly after birth, neonates already display remarkable attunement to their caregivers, demonstrating reflexive and reflex-like patterns of behavior that enable close physical contact [38], as well as distinct patterns of preferential orientation to the sights, sounds, and smells of their caregivers [39,40] (preferences that are present already despite extremely limited *ex utero* experience, with some preferences present already *in utero* [41]). These reflexive behaviors increase an infant's probability of initiating and/or maintaining direct caregiver interaction: the palmar grasp reflex enables holding [38]; the Moro reflex facilitates caregiver contact in the event of falling [42]; the rooting reflex aids neonates' efforts to find mother's nipple [9]; and the sucking reflex facilitates successful and continued feeding [4].

In addition to the presence of these reflexive actions, rather remarkable preferences are evident already at 10 minutes postpartum, when neonates will track a moving face-like pattern farther than a pattern of scrambled or inverted facial features [43]. Similar evidence of preferential orientation has been replicated within the first hour [44] and the first 1–5 days of life [45,46], for cues ranging from biological motion [47] to others' eyes [48]. These preferences are highly specific (Figure 1), encompassing sight [48,49], sound [39], and smell [50,51] (touch, too, undoubtedly plays a critical role [52,53] but remains relatively and unfortunately less well studied).

#### Signaling Behaviors

In order to effectively increase the neonate's chances of survival, the predispositions described above must function as more than mere preference: viewed more broadly, these predispositions must serve not only to direct neonatal resources towards the caregiver, but also to elicit the response of those caregivers by serving as signaling behaviors [54]. Neonatal crying is a clear example of a reflexive behavior that functions as a highly effective signal to promote proximity and contact [55,56]. Crying reliably evokes perceptions of distress and discomfort on the part of most parents [55], is associated with distinct patterns of brain activity, and is believed to evoke distinct neuroendocrine responses (here the existing literature is sparse in humans [57,58] but more well-established in animal models [57]). Crying's efficacy as a signaling behavior is also found in parents' response: holding or cuddling the infant is a highly efficient means of reducing or ending infants' crying [59]. Crying thus signals the infants' state but also elicits parent response and precipitates additional opportunities for interaction.



## Trends in Cognitive Sciences

**Figure 1. Neonatal Preferential Orientation to Caregiver Sounds, Smells, and Sights.** Preferential orienting is observed already in the first days after birth in the auditory domain (A), in response to olfactory cues (B), and in visual attention to faces (C), biological motion (D), and face-like stimuli (E). At 5 days, neonates show distinct preference for looking at faces that have eyes open rather than closed and even distinguish, despite relatively poor visual acuity so soon after birth, between faces with eyes directed towards them rather than away. Similarly, neonates distinguish and prefer their own mother's voice to that of an unknown woman, but prefer the sound of even an unknown woman's voice to that of silence. In addition, neonates recognize their mother's smell and will selectively head-turn at 2–7 days after birth towards the scent of their own mother's breast. Examples of more preferred stimuli are pictured in the left column while less preferred stimuli are pictured in the right column. Stimulus descriptions and citations are given at far right. Data are from references [39,40,43–51,94,132–136,150].

Beyond the vocal domain, the neonate's own facial musculature also serves as a potent signaling system. Fully formed and functional at birth, neonates' facial muscles enable them to produce all but one of the discrete facial actions visible in adults [60]. Although these facial expressions are not initially contingent upon caregiver behavior (e.g., neonatal smiles are largely reflexive, occurring during sleep [61]), the muscle actions themselves clearly resemble interpretable facial expressions, including smiling, brow-knitting, and pouting [60,62], and are fully sufficient to evoke a caregiver's emotional response. By evoking that response (without in any way depending upon it), these noncontingent behaviors facilitate future (actually) contingent social interaction [20,63].

The fact that initially spontaneous, noncontingent actions can beget actual social interaction highlights a means by which adaptive solutions emerge from reoccurring couplings of infant and caregiver [64]. In the preceding example, the capacity of the neonate (the physical ability to form a smiling facial expression) combines with stable, recurrent features of the caregiving environment (the consistency with which the caregiver responds to facial actions that look like smiling). That coupling increases the joint probability of emergent social adaptive action, beginning moments after birth. Critically, the neonate's capacities and the functional significance of those capacities as signaling behaviors cannot be understood without examining them within the caregiving context into which neonates' behaviors are adapted. Through repeated interactions, solutions to the neonate's initial task of survival lead to mutually adaptive actions on the part of both neonate and caregiver.

#### Mutual Adaptation

The behaviors described above (behaviors that serve to preferentially orient neonates' attention towards their caregivers, as well as behaviors that then serve as signals to those caregivers) are adaptive precisely because they provide selective advantage within an environment that includes a caregiver. But those behaviors by themselves would not be sufficient for survival, even in a caregiver's presence. They require the reciprocal, complementary actions of the caregiver, actions that are themselves modified to optimally respond to the signals of the neonate and to then elicit further signals [20].

Evidence of this iterative mutual adaptation (Table 1) is found even upon first contact with the neonate, when mothers display orderly and predictable patterns of species-specific behaviors that include touch and an intense interest in establishing eye-to-eye contact [65]. The mother's interest in the neonate's eyes matches the neonate's ability to attend to mother's eyes [48,49] and appears to play an important role in establishing affective ties [65]. In addition to seeking eye contact, caregivers use slow and exaggerated speech patterns, facial expressions, and body movements that are well matched to the newborn's restricted perceptual capacities [20,66,67]. These are also the actions that appear to optimally engage young infants [66,68].

In another instance of parity between infant and caregiver, newborn crying elicits picking-up and soothing [20], but newborn crying also occurs most frequently when newborns lose visual, auditory, or physical contact with their caregivers [55]. Parental response thus serves to reinstate that which was lost. Picking up the infant not only soothes and promotes state regulation but also increases newborn arousal and visual attentiveness [20,59]. Likewise, touching the infant has the effect of increasing infant eye contact with the caregiver [69]. Each of these mutually adapted actions facilitates still further forms of interaction [70].

Caregivers are, of course, highly motivated to engage their newborns socially. Reflexive actions of the newborn, such as grasping, are often interpreted by caregivers as signals of actual social recognition, affection, or emotional expression, well before such actions are consciously controlled or contingently deployed [20,63]. This perception creates rich opportunities for newborn learning: the meaning that caregivers ascribe to newborns' actions, and the way in which caregivers then modify their own reactions accordingly, creates the ideal framework for newborns to learn the meaning of their own actions: that is, through the reactions of their caregivers [71,72].

Early patterns of imitative behavior reveal this exact form of interactional meaning-making. Mothers imitate their babies' gestures soon after birth and continue to increase their degree of imitation in the following first 3 days [54]. Imitation is a form of mutual adaptation and another

Table 1. Mutual Adaptation of the Infant–Caregiver Dyad

Infant capacity/action	Caregiver action/capacity	Refs
Neonates have strong rooting and sucking reflexes.	Rooting facilitates breastfeeding; sucking triggers maternal prolactin and oxytocin release (prolactin stimulates milk synthesis and oxytocin produces milk release). Amount and frequency of sucking determines the amount of breast milk produced; without suckling, milk production ceases.	[9]
Neonatal sound production is limited (by state of vocal tract development and brain maturation) to crying and vegetative noises (sucking, sneezing, breathing).	Crying reliably elicits caregiver actions, including closer physical proximity, increased holding, and increased cuddling; in lactating women, infant crying stimulates blood flow to the areolar area and milk flow from the breast.	[55,56]
Neonates significantly reduce their crying and become more alert when . . .	caregivers provide proprioceptive-vestibular stimulation (holding and cuddling).	[59]
Neonates demonstrate varying states of alertness.	Caregivers respond to periods of neonate alertness with affection: looking en face, smiling, talking, humming, singing, cuddling, and kissing.	[59]
Infants are differentially sensitive to sounds encompassing and modulated across a broader frequency range (150–550 Hz) than to sounds in narrower, lower frequency ranges (150–275 Hz) or to nonmodulated, lower frequency tones.	Infant-directed speech ('motherese', with characteristically exaggerated pitch contours) covers a broader frequency range inclusive of higher frequency sounds.	[68,146,147]
Newborns demonstrate increased responsiveness to voice sounds.	Newborn responsiveness to voice sounds encourages caregiver interaction.	[20]
Neonates demonstrate differential neural processing of sounds modulated over timescales of 25–50 ms.	Sounds modulated over timescales of 25–50 ms are particularly relevant for spoken communication.	[148,149]
The infant visual system has greater sensitivity to visual signals with high-contrast polarity.	The faces and especially eyes of caregiving conspecifics contain multiple sources of high-contrast visual information.	[94]
The infant visual system has greater sensitivity to low (rather than high) spatial frequency information.	The faces of caregiving conspecifics are replete with low spatial frequency information; facial affect cues are communicated predominantly via low spatial frequencies.	[81]
Infants attend more to top-heavy, vertically asymmetric patterns.	Human faces are vertically asymmetric with features concentrated in their upper portion; mothers display intense interest in establishing eye contact with their infants.	[93]
Neonates spontaneously produce a broad range of facial expressions and gestures.	Spontaneous facial expressions evoke emotional responses in caregivers, and caregivers imitate newborn gestures which, in turn, increases infant gaze towards the caregiver.	[20,60–62]
Infants produce voluntary, intentional facial gesturing.	Caregivers respond to voluntary infant facial gesturing with 'modified imitation': producing contingent responses or producing different but related expressions; caregivers then wait for infant to respond, initiating further cycles.	[54]

strong signaling behavior for both partners. Maternal imitation increases infants' gazing towards the mother, and an infant's reflexive orienting towards the eyes of a caregiver reliably elicits increased caregiver response [54,73].


### Typical Transitions in Social Adaptive Action After 2–3 Months' Experience

Neonates' success in the social adaptive tasks described above (i.e., their success in preferentially directing biological resources towards a caregiver; the success with which their predispositions serve as signaling behaviors; and the success with which those signaling behaviors then become mutually adapted to those of their caregivers) creates, in turn, a set of new conditions that will further facilitate development. The initial acts of orienting, signaling, and

adapting establish myriad opportunities for extended social interaction. These new avenues for social interaction present the infant with new tasks to be learned and put to use, setting the stage for a cascade of transitions in behavior (Figure 2).

### Early Adaptive Action Constrains Later Learning

The iterative process of early infant learning, whereby ‘each [ability] is first of all an effect of the preceding [ability], and itself becomes the cause of the following [ability]’ ([74], p. 112), is well illustrated by the learning that transpires in Rovee-Collier’s classic ‘infant conjugate reinforcement’ paradigm [75]. In this experiment, an infant lies on her back, with a string connecting her foot to a mobile that hangs above. Given the direct coupling between foot and mobile, movements of the infant’s foot induce movement in the mobile, providing the child with novel



Trends in Cognitive Sciences

Figure 2. Developmental Milestones in Infants' First 6 Months. Plotted data, showing the percentage of infants who display each listed behavior between birth and 6 months. Behaviors are plotted sequentially, with behaviors that emerge (and, in some cases, decline) earlier in development at the top and behaviors emerging later in development at the bottom. This sequence of developmental milestones highlights the cascading process of infant development, whereby an infant’s own emerging abilities lead the infant towards new ways of exploring and experiencing the world, prompting further development in an iterative process. Developmental milestone data were scanned or transcribed from published texts and manuals [8,10,12,137–142]; data were fitted with a sigmoid function to describe the cumulative proportion of children displaying each behavior (common milestones were averaged across sources). The fitted functions are color-scaled from generally absent (black) to present in 50% of children (red) to present in most children (yellow).



reinforcement of what were initially only spontaneous leg movements. Importantly, infants discover this relationship themselves: as infants spontaneously kick their feet they learn the contingency between their own movements and the movement of the mobile. Once this link is discovered, the act of kicking, initially spontaneous and unrelated to the mobile, acquires new meaning: kicking becomes an instrumental act for effecting control over the external world. As a consequence of this newfound adaptive value, more precise and vigorous instrumental kicks increase while spontaneous and undirected kicks decline [76].

These data illustrate two key points about developmental change. First, infants play an active role in discovering their own new tasks: in this case, they discover a direct but previously unknown link between their own actions and those of the external world, and they then learn to control and exploit that link. Second, learned volitional behaviors can emerge from and/or depend upon behaviors that were initially spontaneous or reflexive.

Together, these two points demonstrate highly specific constraints that specify and guide subsequent skill acquisition: both the spontaneous initial actions (kicking rather than, for example, hand-waving) as well as the learned solution to that task (kicking with specific magnitude and direction) have the direct effect of limiting and specifying future instances of, and opportunities for, learning. If no initial kicks occurred, or the relationship between kicking and mobile never discerned, the contingency would not be learned. Likewise, by learning to control the mobile with specific kinds of kicks, the probability of other actions (e.g., trying to use hand-waving for the same purpose) declines, while the probability of the newly learned kicking increases (see [77] for an additional elegant example of how early experiences precipitate changes in infants' subsequent activities).


In short: by selectively exploiting specific actions that successfully yielded past adaptive results, infants transform not only their own future behaviors but also their own future possibility of behaviors (i.e., altering the probability that other related behaviors will be more or less likely to occur) [78].

#### Reflex-like Behaviors Give Way to Volitional Actions

The kicking-to-mobile-movement contingency is by no means the only case in which learned, volitional behaviors replace, emerge from, and/or depend upon behaviors that were initially spontaneous or reflexive. A host of such transitions occur in a variety of different developmental domains in an infant's first months (Figure 3).

For instance, the side-to-side head turning reflex, a variant of the rooting reflex, is thought to play an important role in feeding by providing an opportunity for the neonate's mouth to come into contact with the mother's nipple [9]. As neonates gain increasing experience with feeding, this initially spontaneous action transitions to a directed head turning reflex at approximately 2–3 weeks of age [4,9], and then transitions again to volitional directed head turning at approximately 3 months [11]. The hand grasping reflex, present at birth, begins to disappear at approximately 4–5 months of age, coinciding with the emergence of voluntary reaching and grasping [38]. Similarly, the toe grasping reflex, also present at birth, begins to fade as infants gain voluntary control over feet and legs and begin to crawl [10].

These transitions from reflexive behaviors to volitional actions do not appear to progress according to lockstep chronological timers; instead, they depend more heavily on individualized experiences that are necessary to facilitate later transitions. For instance, the directed head turning reflex persists longer in infants who are breast-fed compared with infants who are




Trends in Cognitive Sciences

**Figure 3. Transitions from Reflex-like Actions to Volitional Behaviors.** Behavioral transitions in the first 6 months of life, with implications for early social behavior in autism. (A) Data from references [8,10,12,137–142] showing examples of declining reflex-like actions (unbroken lines) and emerging volitional behaviors (broken lines) in the vocal, visual, and motor domains. Approximate transition times are marked by vertical lines. (B) Top, trajectories of reflex-like actions and volitional behaviors from (A) are plotted against chronological age. Bottom, trajectories are aligned at the time of transition to illustrate the idea that learned, volitional behaviors may replace, emerge from, and/or depend upon behaviors that were initially spontaneous or reflexive: as new volitional behaviors are acquired, the adaptive value of simpler reflex-like actions decreases, as does the action. (C) In typical face perception, existing normative data [44,45,81,92] suggest a similar transition at approximately 2 months: reflex-like eye-looking declines (unbroken gray line) while volitional eye-looking increases (broken gray line). Reflexive eye-looking is believed to be experience-expectant and subcortically mediated, while volitional eye-looking is believed to be experience-dependent and largely cortically mediated. (D) A corresponding reduction in amount of eye-looking by typically developing (TD) infants has been observed at the hypothesized time of transition, shown in the top panel (blue line, sum of percentage fixation on eyes). Observed data from months 2–8 replotted from [32]. Trajectories from 0–2 months represent hypothesized levels of eye-looking in keeping with existing normative data [96,97]. In observed data from [32], infants with autism spectrum disorder (ASD) exhibited relatively high levels of eye-looking at 2 months, which then declined. By contrast, TD infants showed relatively low levels of eye-looking at 2 months, which then increased. Relatively high levels of eye-looking at 2 months in ASD (red) suggests reflex-like eye-looking that is not supplanted by volitional eye-looking and, instead, persists atypically. Rather than an outright failure of cortically controlled voluntary preferential attention in ASD, eye-tracking data suggest a co-opting of experience-dependent cortical mechanisms by attention to other, nonsocial features in the environment. As a result, reflex-like eye-looking gradually declines as it is supplanted by attention to other (non-eye) features. Abbreviation: ATNR, asymmetrical tonic neck reflex.



bottle-fed [4]; voluntary reaching and grasping emerge earlier in infants who are afforded the experience of picking up objects [77]; and the decline of the toe grasping reflex is more closely related to the acquisition of voluntary control of the feet than to chronological age [10]. Each of these examples can be thought of as an individualized timescale of development: initial behaviors provide pivotal opportunities that enable subsequent learning, in highly individualized fashion. (For more on individualized timescales of development, and recent statistical advances in the measurement thereof, see Figure 4.)



Trends in Cognitive Sciences

**Figure 4. Developmental Methods: Quantifying Individualized Timescales of Growth.** Functional data analysis is a relatively new method of analyzing time series data that places greater emphasis on individual trajectories of data, considering the trajectory itself to be a single observation and quantifying that function's variability in terms of both timing and scale [143,144]. This is an important methodological as well as conceptual shift in how longitudinal data are analyzed and understood, with exciting implications for studies of child development and for the conclusions we may draw about underlying biological processes. A good example of this shift can be seen when analyzing a very literal example of a child's growth: change in height. (A) Shows height measurements of girls from 1 to 18 years of age (from the classic Berkeley Growth Study [145]). The underlying biological process of growth is of course nearly identical in all children, but exactly when the pubertal growth spurt occurs and how large it is varies considerably by individual. (B) Individual differences can be expected in timing (when a particular change occurs), in scale (how large or small a given change may be), and in both timing and scale. With conventional growth curve modeling, (C) fitting individual data with a power function (left panels) yields a relatively good fit in statistical terms ( $R^2 > 0.98$  for each of the three example curves); however, it also eradicating all signs of the pubertal growth spurt, as seen especially in the plots of change in height and in rate of change in height (the 1st and 2nd derivatives, respectively). Fitting the same data with a 5th order polynomial [right panels of (C)] improves the picture somewhat, but parameter estimates of when the pubertal spurt occurs in individual children (colored dots) are as much as 2 years earlier than estimates observed in a more data-driven fashion, as in (D), using B-spline basis functions. (D) In functional data analysis, variation in both timing and scale are quantified, and curve shape is determined empirically. (E) Rather than being confounded by individual differences in maturational rate (individualized developmental timescale), functional data analysis measures the extent of these differences as 'warping' or registration functions, explicitly comparing and correcting for differences in chronological time versus individual maturational time. (F) When data are analyzed as functional trajectories, registered according to measures of individual difference in developmental timing, the ability to estimate the shape of the actual developmental process improves substantially.

### Infant–Caregiver Adaptation Guides Developmental Transitions

The developmental learning processes described above are equally applicable to transitions in social learning. Just as infants discover a physical linkage in the foot and mobile example, infants also discover links connecting them to their social partners [79,80]. Research on infant–caregiver dyads reveals that infants' facial movements, head movements, and vocalizations are coupled, often tightly, to the facial expressions and sounds of their caregivers [54,79]. As infants discover and exploit the links between their own actions and the actions and reactions of others, the spontaneous and reflex-like behaviors of the neonate are transformed.

One notable example of such a transition is the case of orienting towards faces. Preferential tracking of faces (as described above) is initially a reflex-like, spontaneous, stimulus-driven form of orienting, observed from the first hours after birth [43]. Notably, however, this neonatal orienting response declines after 4–6 weeks postpartum, reaching a relative low point, before preferential attention to faces then increases at approximately 2 months [29,81]. This would appear to be a transition from a more reflex-like response (i.e., not socially contingent) to a more volitional, truly social, form of orienting. Evidence of this transition is also supported by other changes at approximately 2 months. First, the physical characteristics of face-like stimuli that most effectively elicit preferential attention also change: while neonatal preference for faces can be elicited by simple schematic stimuli [44], real faces are preferred by 1–2-month-old and older infants (reviewed in [82,83]).

In addition, the 2nd month also marks the first time in which clear evidence exists that infants engage with other people in an interactive, socially meaningful manner. At approximately 2 months, infants begin smiling and cooing in response to faces [84]. Prior to this point, neonatal smiling is endogenous, happens most often during sleep, and shows little evidence of linkage to external stimulation [61,84]. At 2 months, however, smiles begin to be contingently linked to changes in caregiver vocalization and affective state, indicating not only the emergence of social smiling but also indicating that infants' preferential visual attention (in this case, directed towards faces) now also functions as a form of communication [84]. Newborn vocal development also undergoes marked change shortly thereafter: prior to approximately 3 months, crying is reflexive, undifferentiated, and marked by an absence of vowel production; after 3 months, cries become interactive, intentional [85], and increasingly reflective of the infant's psychophysiological state [3] (note that the timing of these changes, in contrast to smiling, is partially constrained by anatomy: at 3 months, the larynx descends [3] and the ribs reconfigure [86], altering the infant's capacity for vocal production).

Finally, there are also large changes at approximately 2 months in infants' expectations of other people. New capacities for active social engagement are evidenced by several classic studies: in the still-face paradigm [87], when a formerly interactive partner becomes unresponsive, 1.5- and 3-month-old infants (but not neonates) become upset, with signs of distress including crying, general negative affect, and shifts in visual attention away from the caregiver [88]. In addition, infant social bids during the still-face phase increase at 2 months of age [80], offering evidence that infants have now become active agents in dyadic interaction, developing specific expectations of the ways in which social interactions should unfold.

As infants exploit the links between their own actions and those of their caregivers, caregivers, in turn, modify their behavior to progressively advance cycles of learning. During the infant's second month, mothers spend more time talking and smiling with their infants, and less time displaying neutral facial expressions [89]. Regularities in face-to-face play (such as turn-taking) are established, repeated, and then extended during this period [54], with the peak in face-to-

face play occurring between 3 and 6 months [90]. Evidence suggests that mothers of 3-month-olds display an average of approximately eight changes in emotion-related facial expressions per minute [54]. With even 3 hours per day of direct interaction, and with as little as 25% of that time spent directly attending to mother's face [91], an infant would be exposed to approximately 360 changing emotional cues in a single day, or more than 32 000 between the ages of 3 and 6 months.

#### Co-occurring Changes in Early Infant Brain and Body Systems

Relative to our knowledge of infant behavioral transitions, detailed information about co-occurring changes in brain are still limited. Existing knowledge does, however, suggest a substantial transition also occurring at approximately 2 months, shifting from largely subcortical to increasingly cortical control [92].


At birth, reflexive visual orienting appears to be supported by the subcortical visual system, which is differentially responsive to low-spatial frequency information [81], top-heavy vertically asymmetric patterns [93], and high contrast polarity [94]. Notably, these are all properties also strongly represented in the faces and eyes of conspecifics, and much like other mutual adaptations between infant and caregiver (Table 1), this match between signal and receiver may be a means of increasing the probability that neonates orient towards conspecifics rather than other competing stimuli [81,95].

As the neonate enters her second postnatal month, existing models suggest that reflexive orienting responses, subserved by subcortical structures, decline due to developing inhibition by cortical circuits (as visual preferences come under increasing cortical control) [96]. This theory is supported by the relative immaturity of primary visual cortex at birth [97] and also by developmental changes in orienting. Newborn orienting is most effectively elicited by face-like stimuli presented in the peripheral visual field [45] (the area that feeds more directly into subcortical visual pathways [98]), whereas preferential orienting in 2-month-olds is more effectively elicited by stimuli presented in the central visual field (which feeds more directly into cortical visual pathways [97]). Further evidence comes from the fact that retinocortical pathways become fully functional at approximately 2 months [99], and signs of adult-like cortical specialization for face processing are also first observed at 2- to 3-months in both event-related potential (ERP) [100] and positron emission tomography studies [101].

Generally speaking, however, much is still to be learned about infant brain development and the manner in which behavioral milestones are coupled with brain changes. Newly available resources (e.g., the Allen Human Brain Atlas [102] and BrainSpan [103,104] projects) and ongoing projects (Human Connectome [105] and Developing Human Connectome [106] Projects) are likely to transform our understanding of infant brain development in the coming years (Figure 5). Currently, however, data from infant behavior and brain development showcase a system that undergoes enormous developmental change in a matter of months: neonatal preferences and reflex-like responding give way to remarkably sensitive and contingent volitional action. We now review some of the emerging literature regarding what is currently known about the derailment of such action in children with ASD.

#### Disruptions in Social Adaptive Action in Autism Spectrum Disorder

A striking developmental contrast in autism is that older children, and even adults, on the autism spectrum often exhibit deficits in social adaptive action that are otherwise present in typical infancy. For example, while typically developing infants preferentially orient to the eyes of others [48,49], 2-year-olds [107], 10-year-olds [108], and even adults [109] with ASD do not.



Trends in Cognitive Sciences

**Figure 5. Longitudinal Expression of Genes Associated With Neurodevelopmental Processes.** Recent advances in studying the spatio-temporal dynamics of gene expression in the human brain augur a new frontier for studies of infant and child development, connecting well-studied behavioral and cognitive milestones to multiple measures of infant brain biology. One example is the BrainSpan Atlas of the Developing Human Brain<sup>1</sup> [104], offering a transcriptional architecture of the human brain from early fetal development through adulthood. While longitudinal expression patterns for more than 17,500 genes were analyzed, the gene expression trajectories plotted here are for genes whose differential expression is associated with key neurodevelopmental processes. (A) Fourteen brain regions in which expression was measured are highlighted. (B) Five sets of genes are highlighted, the expression levels of which are associated with neurogenesis, synaptogenesis, dendrite development, axon development, and myelination, respectively. (C) Longitudinal expression levels of genes associated with each process are plotted, for each brain region, as a percentage of the minimum and maximum lifetime levels of expression. Brain regions are sorted according to peak expression levels between 0 and 24 postnatal months of age. Note the clear waves of expression during these periods of early child development: high levels of expression associated with neurogenesis continue through the first 6 months of life, with marked decline thereafter, followed by waves of increased expression associated with synaptogenesis, dendrite development, axon development, and finally myelination. Given the enormous developmental change occurring in these time periods, data like these, coupled with densely sampled behavioral and neuroimaging data, are likely to transform understanding of infant development in the coming years. See online supplemental materials for additional related references.

Unfortunately, direct observation of infant development in autism is still limited. Given the late average age of conventional diagnosis [110], and the extent to which diagnosis currently depends upon behaviors that emerge in toddlerhood [111], a large gap in current knowledge remains. To fill that gap, the past 10 years have seen a substantial increase in prospective

studies of the infant siblings of children with ASD, whose risk of also having the condition is approximately 20% [112], a substantial increase relative to infants in the general population [113].

#### Social Adaptive Action and Brain Specialization in Autism Spectrum Disorder

Recent studies examining the course of early social development in high-risk infant siblings have focused primarily on measures of social attention, interaction, and/or more general aspects of social development, using standardized assessments as well as experimental tasks (see online supplemental materials for references). A wide array of between-group differences have been identified among infants subsequently diagnosed with ASD, including, among others, differences in frequency of shared smiles, vocalizations, and gaze to faces [114,115]; attention to social scenes and images [116]; attention to and engagement with objects [117]; ERP amplitude differences in relation to observed eye gaze [118]; temperament profiles [119]; responsiveness to distress in others [120]; mutuality of infant-caregiver interaction [121]; and social smiling and joint attention [122,123].

Related differences are also found in brain structure [34,124,125] and function [33], with differences in extra-axial cerebrospinal fluid volume [125], cortical surface area [34], white matter fiber tracts [124], and functional connectivity [33] all identified as possible predictors of autism in the first year of life. Some studies have also highlighted ways in which similar overt behaviors were accompanied by notable differences in measures of underlying brain function [118,126].

Across these varied studies, one relatively consistent finding is that although indications of earlier differences are present, between-group differences are more easily identified at later ages (see online supplemental materials for references). This pattern of results is consistent with the notion reviewed above that development is iterative, with an infant's own actions and experiences guiding and constraining future learning [78]. Such a model predicts that even small disruptions of early social adaptive action can yield progressively greater between-group differences over time, culminating later in the marked symptomatology of ASD. Importantly, recognition of development as an iterative process means that differences observed at later times must be underway already at earlier ones.

#### Early Patterns of Preferential Orientation in Autism Spectrum Disorder and Transitions Therein

Given the foundational nature of social skills disruption in ASD, as well as the profound transitions in typical social development happening in early infancy, changes in preferential orientation, engagement, and adaptation during the first weeks and months after birth are a critical area of future ASD inquiry.

Existing data in infants later diagnosed with ASD suggest that one such mechanism, preferential orientation to the eyes, is not immediately diminished [32]. Levels of early eye-looking are relatively high in infants with ASD. To be clear, the presence of these relatively high levels does not necessarily indicate 'normative' or 'intact' function [32]; to the contrary, while the magnitude of eye-looking at 2 months is relatively high, the underlying developmental processes are clearly already different: rate-of-change in eye-looking, from at least 2 months onwards, differs significantly between outcome groups. Moreover, infants later diagnosed with ASD exhibit a slight but statistically significant increase in eye-looking at 2 months, which then declines (Figure 3D); by contrast, typically developing infants exhibit a relative low point in eye-looking at 2 months, which then increases.

As reviewed above, the timing of these differences may relate directly to important transitions in adaptive behavior: preferential orientation to faces (present at birth [43]) typically declines between 4 and 6 weeks [44] before beginning to increase at approximately 2 months [81]. Similar transitions during this same period are observed in orienting to auditory sounds [127] and imitating others [128]. At a neural level, these changes are thought to be accompanied by a shift from subcortical to cortical control, with initial reflex-like predispositions (subserved by subcortical structures) declining as cortical control matures [81].

Taken as a whole, a very specific hypothesis emerges (Figure 3D): reflex-like orienting to the eyes of others may be initially, at least superficially, present in newborns with ASD, while the latter phase of preferential eye-looking is disrupted. More broadly, in a wide range of other sensory and behavioral domains, preferential orientation may initially be present in these infants, whilst early pivotal transitions (transitions that depend upon the way in which initial preferential orientation is integrated into mutually adaptive interaction) are disrupted. Critically, this hypothesis focuses scientific attention on key developmental transitions very early in life, and specifically on the progression from a reflexive/‘experience expectant’ phase of development (largely subcortically mediated), to an interactive/‘experience dependent’ phase (with greater dependence on cortical control) [96,97].

To be clear, there is little evidence to suggest an outright failure of cortical control of voluntary preferential visual attention in ASD; to the contrary, data at later age points suggest a co-opting of those cortical mechanisms by attention to other features of the environment, such as nonsocial, physical cues [107,129]. These data suggest future lines of inquiry, focused in these first few months, on how early predispositions transition into mutually adapted social interaction. This line of inquiry should benefit from a direct link to decades of work from pertinent model systems [130], work that can now be brought to bear in guiding gene–brain–behavior hypotheses of the pathogenesis of autism. Likewise, the role of other sensory systems in these processes remains understudied; given the early development of the auditory and tactile systems (affective touch in particular [53]), their role in constraining and influencing early development is a key area for future study.

### Concluding Remarks

Collectively, the data reviewed above, from both behavior and brain development in typical infancy, showcase a system that is remarkably attuned to the social world, is mutually adapted to caregiver behavior, and becomes progressively more so over time. From the first moments of life, neonates exhibit a wide range of socially adaptive preferences and reflex-like responses that serve to orient their attention towards caregivers as well as behaviors that serve as important signals to those caregivers. These behaviors are adaptive because they occur within an environment that depends upon a caregiver, one whose own actions are adapted to optimally respond to the neonate’s and to further elicit such actions. This mutually adapted, mutually reinforcing context of the infant–caregiver dyad is both the platform and the catalyst for subsequent development: initially spontaneous reflex-like responses transition into remarkably sensitive and contingent social action, all within the first months of life.

These findings, together with evidence from early normative transitions, provide an important framework for future research in ASD by constraining hypotheses about what specific developmental transitions may be disrupted in ASD and when. For instance, rather than suggesting a complete absence of preferential orientation to others from birth (as was hypothesized dating back to Kanner [131]), these data identify a specific transitional period of early infancy, around the second month of life, as a critical focus for future investigation, both for systems level

### Outstanding Questions

How do neonatal reflexive actions and predispositions transition into mutually adapted social interaction in the first months of typical development? What changes at the level of brain networks and gene expression accompany such transitions?

How do infants’ early behaviors facilitate later attainment of normative social developmental milestones?

How do earlier-developing systems (e.g., touch), as well as disruptions in such systems (e.g., potential disruptions in affective touch in infants with ASD), have the potential to constrain and canalize the development of later-emerging systems?

For children with delays in specific milestones, is there a corresponding delay in offset of related reflexes? Or, alternatively, do reflexes decline on their own timeline, irrespective of other subsequent delays?

Is the transition from ‘experience expectant’ reflexive action to ‘experience dependent’ social interaction accompanied by a shift from subcortical to cortical control?

Are typical predispositions for conspecific orienting present in newborns with ASD?

How and when do mechanisms of preferential orientation, signaling, and infant–caregiver adaptation become disrupted in infants with ASD?



neuroscience research (in both human and model systems) and for genetics research (guiding and providing the quantitative traits for developmental gene expression and methylation studies). These data further suggest that future analysis of potential disruptions in autism might productively focus on mechanisms of preferential orientation, signaling, and, critically, mutual adaptation between infant and caregiver, particularly during this transitional period (see Outstanding Questions). The pursuit of these constrained hypotheses holds great promise for yielding insight into the specific developmental processes that are disrupted in ASD and for discovering new opportunities for innovative early interventions.

### Acknowledgments

This work was supported by grants from the National Institute of Mental Health: NIMH R21 MH105816; NIMH P50 MH100029; NIMH 2P50 MH100029-6; and NIMH K01 MH108741. Additional support provided by the Marcus Foundation, the Whitehead Foundation, the Simons Foundation, and the Georgia Research Alliance. We would also like to thank Torrey Cohenour, Aiden Ford, Drew Kreuzman, Sarah Markert, Megan Micheletti, and Jack Olmstead, for helpful comments on the manuscript, as well as Emily Ryu for assistance in gathering infant developmental milestone data.

### Resources

[www.brainspan.org](http://www.brainspan.org)

### Appendix A Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.tics.2018.02.012>.

### References

- Nagy, E. (2011) The newborn infant: a missing stage in developmental psychology. *Infant Child Dev.* 20, 3–19
- Wolff, P.H. and Ferber, R. (1979) The development of behavior in human infants, premature and newborn. *Annu. Rev. Neurosci.* 2, 291–307
- Lieberman, P. (1985) The physiology of cry and speech in relation to linguistic behavior. In *Infant Crying* (Lester, B.M., ed.), pp. 29–57, Plenum Press
- Ingram, T.T.S. (1962) Clinical significance of the infantile feeding reflexes. *Dev. Med. Child Neurol.* 4, 159–169
- Delaney, A.L. and Arvedson, J.C. (2008) Development of swallowing and feeding: prenatal through first year of life. *Dev. Disabil. Res. Rev.* 14, 105–117
- Parmelee, A.H. and Stern, E. et al. (1972) Development of states in infants. In *Sleep and the Maturing Nervous System* (Clemente, C., ed.), pp. 199–228, Academic Press
- Bronson, G. (1974) The postnatal growth of visual capacity. *Child Dev.* 45, 873–890
- Gingold, M.K. et al. (1998) The rise and fall of the plantar response in infancy. *J. Pediatr.* 133, 568–570
- Prechtl, H.F.R. (1958) The directed head turning response and allied movements of the human baby. *Behaviour* 13, 212–242
- Gentry, E.F. and Aldrich, C.A. (1948) Toe reflexes in infancy and the development of voluntary control. *Am. J. Dis. Child.* 76, 389–400
- Papoušek, H. (1961) Conditioned head rotation reflexes in infants in the first months of life. *Acta Paediatr.* 50, 565–576
- Twitchell, T.E. (1965) The automatic grasping responses of infants. *Neuropsychologia* 3, 247–259
- Holland, D. et al. (2014) Structural growth trajectories and rates of change in the first 3 months of infant brain development. *JAMA Neurol.* 71, 1266–1274
- Gilmore, J.H. et al. (2007) Regional gray matter growth, sexual dimorphism, and cerebral asymmetry in the neonatal brain. *J. Neurosci.* 27, 1255–1260
- Groeschel, S. et al. (2010) Developmental changes in cerebral grey and white matter volume from infancy to adulthood. *Int. J. Dev. Neurosci.* 28, 481–489
- Huttenlocher, P.R. (1979) Synaptic density in human frontal cortex—developmental changes and effects of aging. *Brain Res.* 163, 195–205
- Huttenlocher, P.R. (1990) Morphometric study of human cerebral cortex development. *Neuropsychologia* 28, 517–527
- Petanjek, Z. et al. (2011) Extraordinary neoteny of synaptic spines in the human prefrontal cortex. *Proc. Natl. Acad. Sci. U. S. A.* 108, 13281–13286
- Gibson, E.J. (1988) Exploratory behavior in the development of perceiving, acting, and the acquiring of knowledge. *Annu. Rev. Psychol.* 39, 1–41
- Lozoff, B. et al. (1977) The mother-newborn relationship: limits of adaptability. *J. Pediatr.* 91, 1–12
- Bowlby, J. (1969) *Attachment and Loss*, Basic Books
- Stern, D.N. (1974) Mother and infant at play: the dyadic interaction involving facial, vocal, and gaze behaviors. In *The Effect of the Infant on its Caregivers* (Lewis, M. and Rosenblum, L.A., eds), Wiley-Interscience
- Brazelton, T.B. et al. (1975) Early mother-infant reciprocity. *Ciba Found. Symp.* 33, 137–154
- Snow, C.E. (1977) The development of conversation between mothers and babies. *J. Child Lang.* 4, 1–22
- Papoušek, M. et al. (1995) Origins of reciprocity and mutuality in prelinguistic parent-infant “dialogues”. In *Mutualities in Dialogue* (Marková, I., ed.), pp. 58–81, Cambridge University Press
- Tronick, E.Z. (1995) Touch in mother-infant interaction. In *Touch in Early Development* (Field, T., ed.), pp. 53–65, Lawrence Erlbaum Associates
- Hertenstein, M.J. (2002) Touch: its communicative functions in infancy. *Hum. Dev.* 45, 70–94
- Ainsworth, M.D.S. and Bell, S.M. (1974) Mother-infant interaction and the development of competence. In *The Growth of Competence* (Connolly, K.J. and Bruner, J.S., eds), Academic Press

29. Feldman, R. (2015) The adaptive human parental brain: implications for children's social development. *Trends Neurosci.* 38, 387–399
30. Stern, D.N. (2009) *The First Relationship: Infant and Mother*, Harvard University Press
31. Hobson, P. (2004) *The Cradle of Thought: Exploring the Origins of Thinking*, Pan Macmillan
32. Jones, W. and Klin, A. (2013) Attention to eyes is present but in decline in 2-6-month-old infants later diagnosed with autism. *Nature* 504, 427–431
33. Emerson, R.W. et al. (2017) Functional neuroimaging of high-risk 6-month-old infants predicts a diagnosis of autism at 24 months of age. *Sci. Transl. Med.* 9, 1–8
34. Hazlett, H.C. et al. (2017) Early brain development in infants at high risk for autism spectrum disorder. *Nature* 542, 348–351
35. Poirier, F.E. and Smith, E.O. (1974) Socializing functions of primate play. *Integr. Comp. Biol.* 14, 275–287
36. Bruner, J.S. (1972) Nature and uses of immaturity. *Am. Psychol.* 27, 687
37. Greenough, W.T. et al. (1987) Experience and brain development. *Child Dev.* 58, 539–559
38. Gesell, A. and Ámatruda, C.S. (1947) *Developmental Diagnosis*. (2nd edn), Hoeber
39. DeCasper, A.J. and Fifer, W.P. (1980) Of human bonding: newborns prefer their mothers' voices. *Science* 208, 1174–1176
40. Bushnell, I.W.R. et al. (1989) Neonatal recognition of the mother's face. *Br. J. Dev. Psychol.* 7, 3–15
41. Kisilevsky, B.S. et al. (2003) Effects of experience on fetal voice recognition. *Psychol. Sci.* 14, 220–224
42. Moro, E. (1918) Das erste trimenon. *Münchener Med. Wochenschr.* 65, 1147–1150
43. Goren, C.C. et al. (1975) Visual following and pattern discrimination of face-like stimuli by newborn infants. *Pediatrics* 56, 544–549
44. Johnson, M.H. et al. (1991) Newborns' preferential tracking of face-like stimuli and its subsequent decline. *Cognition* 40, 1–19
45. Simion, F. et al. (1998) Preferential orienting to faces in newborns: a temporal–nasal asymmetry. *J. Exp. Psychol. Hum. Percept. Perform.* 24, 1399
46. Valenza, E. et al. (1996) Face preference at birth. *J. Exp. Psychol. Hum. Percept. Perform.* 22, 892
47. Simion, F. et al. (2008) A predisposition for biological motion in the newborn baby. *Proc. Natl. Acad. Sci. U. S. A.* 105, 809–813
48. Farroni, T. et al. (2002) Eye contact detection in humans from birth. *Proc. Natl. Acad. Sci. U. S. A.* 99, 9602–9605
49. Batki, A. et al. (2000) Is there an innate gaze module? Evidence from human neonates. *Infant Behav. Dev.* 23, 223–229
50. Porter, R.H. and Winberg, J. (1999) Unique salience of maternal breast odors for newborn infants. *Neurosci. Biobehav. Rev.* 23, 439–449
51. Macfarlane, A. (1975) Olfaction in the development of social preferences in the human neonate. *Ciba Found. Symp.* 33, 103–117
52. Field, T. (2002) Infants' need for touch. *Hum. Dev.* 45, 100–103
53. McGlone, F. et al. (2014) Discriminative and affective touch: sensing and feeling. *Neuron* 82, 737–755
54. Malatesta, C.Z. and Izard, C.E. (1984) The ontogenesis of human social signals: from biological imperative to symbol utilization. In *The Psychobiology of Affective Development* (Fox, N. A. and Davidson, R.J., eds), pp. 161–206, Erlbaum
55. Bell, S.M. and Ainsworth, M.D. (1972) Infant crying and maternal responsiveness. *Child Dev.* 43, 1171–1190
56. Soltis, J. (2004) The signal functions of early infant crying. *Behav. Brain Sci.* 27, 443–458
57. Newman, J.D. (2007) Neural circuits underlying crying and cry responding in mammals. *Behav. Brain Res.* 182, 155–165
58. Swain, J.E. et al. (2011) Neuroendocrinology of parental response to baby-cry. *J. Neuroendocrinol.* 23, 1036–1041
59. Korner, A.F. and Grobstein, R. (1966) Visual alertness as related to soothing in neonates: implications for maternal stimulation and early deprivation. *Child Dev.* 37, 867–876
60. Oster, H. and Ekman, P. (1978) Facial behavior in child development. In *Minnesota Symposia on Child Psychology*, (Vol. 11), pp. 231–276, Lawrence Erlbaum Associates
61. Messinger, D. et al. (2002) How sleeping neonates smile. *Dev. Sci.* 5, 48–54
62. Oster, H. (1978) Facial expression and affect development. In *The Development of Affect* (Lewis, M. and Rosenblum, L.A., eds), pp. 43–75, Springer
63. Greenberg, M. and Morris, N. (1974) Engrossment: the newborn's impact upon the father. *Am. J. Orthopsychiatry* 44, 520
64. Thelen, E. and Smith, L.B. (1998) Dynamic systems theories. In *Handbook of Child Psychology* (Damon, W., ed.), pp. 563–635, Wiley
65. Klaus, M.H. et al. (1970) Human maternal behavior at the first contact with her young. *Pediatrics* 46, 187–192
66. Brand, R.J. and Shallock, W.L. (2008) Infants prefer motionless to adult-directed action. *Dev. Sci.* 11, 853–861
67. Fernald, A. et al. (1992) Human maternal vocalizations to infants as biologically relevant signals: an evolutionary perspective. In *The Adapted Mind: Evolutionary Psychology and the Generation of Culture* (Barkow, J., ed.), pp. 391–428, Oxford University Press
68. Cooper, R.P. and Aslin, R.N. (1990) Preference for infant-directed speech in the first month after birth. *Child Dev.* 61, 1584–1595
69. Peláez-Nogueras, M. et al. (1996) Infants' preference for touch stimulation in face-to-face interactions. *J. Appl. Dev. Psychol.* 17, 199–213
70. Nakayama, H. (2015) Emergence of amae crying in early infancy as a possible social communication tool between infants and mothers. *Infant Behav. Dev.* 40, 122–130
71. Klin, A. (1989) Understanding early infantile autism: an application of GH Mead's theory of the emergence of mind. *L. S. E. Quarterly* 3, 336–356
72. Mead, G.H. (1934) *Mind, Self & Society: From the Standpoint of a Social Behaviorist*, The University of Chicago Press
73. Field, T.M. (1977) Effects of early separation, interactive deficits, and experimental manipulations on infant-mother face-to-face interaction. *Child Dev.* 48, 763–771
74. Hall, B.K. (1999) *Evolutionary Developmental Biology*, Springer Science & Business Media
75. Rovee, C.K. and Rovee, D.T. (1969) Conjugate reinforcement of infant exploratory behavior. *J. Exp. Child Psychol.* 8, 33–39
76. Thelen, E. and Fisher, D.M. (1983) From spontaneous to instrumental behavior: kinematic analysis of movement changes during very early learning. *Child Dev.* 54, 129–140
77. Needham, A. et al. (2002) A pick-me-up for infants' exploratory skills: early simulated experiences reaching for objects using "sticky mittens" enhances young infants' object exploration skills. *Infant Behav. Dev.* 25, 279–295
78. Gibson, E.J. and Pick, A.D. (2000) *An Ecological Approach to Perceptual Learning and Development*, Oxford University Press
79. Smith, L.B. (2005) Cognition as a dynamic system: principles from embodiment. *Dev. Rev.* 25, 278–298
80. Bigelow, A.E. and Power, M. (2016) Effect of maternal responsiveness on young infants' social bidding-like behavior during the still face task. *Infant Child Dev.* 25, 256–276
81. Morton, J. and Johnson, M.H. (1991) CONSPEC and CONLERN: a two-process theory of infant face recognition. *Psychol. Rev.* 98, 164–181
82. Sherrod, L.R. (1981) Issues in cognitive-perceptual development: the special case of social stimuli. In *Infant Social Cognition: Empirical and Theoretical Considerations*, pp. 11–36, Lawrence Erlbaum Associates

83. Langsdorf, P. *et al.* (1983) Interest expression, visual fixation, and heart rate changes in 2- and 8-month-old infants. *Dev. Psychol.* 19, 375–386
84. Messinger, D. and Fogel, A. (2007) The interactive development of social smiling. *Adv. Child Dev. Behav.* 35, 327–366
85. Ostwald, P. and Murray, T. (1985) The communicative and diagnostic significance of infant sounds. In *Infant Crying: Theoretical and Research Perspectives* (Lester, B. and Boukydis, C. F.Z., eds), pp. 139–158, Plenum Press
86. Langlois, A. *et al.* (1980) Pre-speech respiratory behavior during the first year of life. In *Infant Communication: Cry and Early Speech* (Murry, T. and Murry, J., eds), College-Hill
87. Tronick, E. *et al.* (1979) The infant's response to entrapment between contradictory messages in face-to-face interaction. *J. Am. Acad. Child Psychiatry* 17, 1–13
88. Bertin, E. and Striano, T. (2006) The still-face response in newborn, 1.5-, and 3-month-old infants. *Infant Behav. Dev.* 29, 294–297
89. Lavelli, M. and Fogel, A. (2005) Developmental changes in the relationship between the infant's attention and emotion during early face-to-face communication: the 2-month transition. *Dev. Psychol.* 41, 265–280
90. Lamb, M.E. *et al.* (1987) The development of infant social expectations in face-to-face interaction: a longitudinal study. *Merrill Palmer Q. (Wayne State Univ. Press)* 33, 241–254
91. Hittelman, J. and Dicks, R. (1979) Sex difference in neonatal eye contact time. *Merrill Palmer Q. (Wayne State Univ. Press)* 24, 171–184
92. Johnson, M.H. *et al.* (2015) The two-process theory of face processing: modifications based on two decades of data from infants and adults. *Neurosci. Biobehav. Rev.* 50, 169–179
93. Simion, F. *et al.* (2001) The origins of face perception: specific versus non-specific mechanisms. *Infant Child Dev.* 10, 59–65
94. Faroni, T. *et al.* (2005) Newborns' preference for face-relevant stimuli: effects of contrast polarity. *Proc. Natl. Acad. Sci. U. S. A.* 102, 17245–17250
95. Johnson, M.H. (2005) Subcortical face processing. *Nat. Rev. Neurosci.* 6, 766–774
96. Johnson, M.H. and Morton, J. (1991) *Biology and Cognitive Development: The Case of Face Recognition*, Blackwell Scientific Publications
97. Johnson, M. (1990) Cortical maturation and the development of visual attention in early infancy. *J. Cogn. Neurosci.* 2, 81–95
98. Rafal, R. *et al.* (1991) Extrageniculate contributions to reflex visual orienting in normal humans: a temporal hemifield advantage. *J. Cogn. Neurosci.* 3, 322–328
99. Braddick, O.J. *et al.* (1986) Orientation-specific cortical responses develop in early infancy. *Nature* 320, 617–619
100. Halit, H. *et al.* (2003) Cortical specialisation for face processing: face-sensitive event-related potential components in 3- and 12-month-old infants. *Neuroimage* 19, 1180–1193
101. Tzourio-Mazoyer, N. *et al.* (2002) Neural correlates of woman face processing by 2-month-old infants. *Neuroimage* 15, 454–461
102. Hawrylycz, M.J. *et al.* (2012) An anatomically comprehensive atlas of the adult human brain transcriptome. *Nature* 489, 391–399
103. Miller, J.A. *et al.* (2014) Transcriptional landscape of the prenatal human brain. *Nature* 508, 199–206
104. Kang, H.J. *et al.* (2011) Spatio-temporal transcriptome of the human brain. *Nature* 478, 483–489
105. Van Essen, D.C. *et al.* (2013) The WU-Minn Human Connectome Project: an overview. *Neuroimage* 80, 62–79
106. Hughes, E.J. *et al.* (2017) A dedicated neonatal brain imaging system. *Magn. Reson. Med.* 78, 794–804
107. Jones, W. *et al.* (2008) Absence of preferential looking to the eyes of approaching adults predicts level of social disability in 2-year-old toddlers with autism spectrum disorder. *Arch. Gen. Psychiatry* 65, 946–954
108. Rice, K. *et al.* (2012) Parsing heterogeneity in autism spectrum disorders: visual scanning of dynamic social scenes in school-aged children. *J. Am. Acad. Child Adolesc. Psychiatry* 51, 238–248
109. Klin, A. *et al.* (2002) Visual fixation patterns during viewing of naturalistic social situations as predictors of social competence in individuals with autism. *Arch. Gen. Psychiatry* 59, 809–816
110. Shattuck, P.T. *et al.* (2009) Timing of identification among children with an autism spectrum disorder: findings from a population-based surveillance study. *J. Am. Acad. Child Adolesc. Psychiatry* 48, 474–483
111. American Psychiatric Association (2013) *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*, American Psychiatric Association
112. Ozonoff, S. *et al.* (2011) Recurrence risk for autism spectrum disorders: a baby siblings research consortium study. *Pediatrics* 128, e488–e495
113. Wingate, M. *et al.* (2014) Prevalence of autism spectrum disorder among children aged 8 years-autism and developmental disabilities monitoring network, 11 sites, United States, 2010. *MMWR Surveill. Summ.* 63, 1–21
114. Ozonoff, S. *et al.* (2010) A prospective study of the emergence of early behavioral signs of autism. *J. Am. Acad. Child Adolesc. Psychiatry* 49, 256–266
115. Filliter, J.H. *et al.* (2015) Positive affect in infant siblings of children diagnosed with autism spectrum disorder. *J. Abnorm. Child Psychol.* 43, 567–575
116. Chawarska, K. *et al.* (2013) Decreased spontaneous attention to social scenes in 6-month-old infants later diagnosed with autism spectrum disorders. *Biol. Psychiatry* 74, 195–203
117. Elsabbagh, M. *et al.* (2013) Disengagement of visual attention in infancy is associated with emerging autism in toddlerhood. *Biol. Psychiatry* 74, 189–194
118. Elsabbagh, M. *et al.* (2012) Infant neural sensitivity to dynamic eye gaze is associated with later emerging autism. *Curr. Biol.* 22, 338–342
119. Garon, N. *et al.* (2016) Temperament and its association with autism symptoms in a high-risk population. *J. Abnorm. Child Psychol.* 44, 757–769
120. Campbell, S.B. *et al.* (2015) Concern for another's distress in toddlers at high and low genetic risk for autism spectrum disorder. *J. Autism Dev. Disord.* 45, 3594–3605
121. Campbell, S.B. *et al.* (2015) Social engagement with parents in 11-month-old siblings at high and low genetic risk for autism spectrum disorder. *Autism* 19, 915–924
122. Sullivan, M. *et al.* (2007) Response to joint attention in toddlers at risk for autism spectrum disorder: a prospective study. *J. Autism Dev. Disord.* 37, 37–48
123. Lambert-Brown, B.L. *et al.* (2015) Positive emotional engagement and autism risk. *Dev. Psychol.* 51, 848–855
124. Wolff, J.J. *et al.* (2012) Differences in white matter fiber tract development present from 6 to 24 months in infants with autism. *Am. J. Psychiatry* 169, 589–600
125. Shen, M.D. *et al.* (2017) Increased extra-axial cerebrospinal fluid in high-risk infants who later develop autism. *Biol. Psychiatry* 82, 186–193
126. Key, A.P.F. and Stone, W.L. (2012) Same but different: 9-month-old infants at average and high risk for autism look at the same facial features but process them using different brain mechanisms. *Autism Res.* 5, 253–266
127. Field, J. *et al.* (1980) Infants' orientation to lateral sounds from birth to three months. *Child Dev.* 51, 295–298
128. Bjorklund, D. (1987) A note on neonatal imitation. *Dev. Rev.* 7, 86–92
129. Klin, A. *et al.* (2009) Two-year-olds with autism orient to non-social contingencies rather than biological motion. *Nature* 459, 257–261
130. Rosa Salva, O. *et al.* (2011) The evolution of social orienting: evidence from chicks (*Gallus gallus*) and human newborns. *PLoS One* 6, e18802

131. Kanner, L. (1943) Autistic disturbances of affective contact. *Nerv. Child* 2, 217–250
132. Vouloumanos, A. and Werker, J.F. (2007) Listening to language at birth: evidence for a bias for speech in neonates. *Dev. Sci.* 10, 159–164
133. Butterfield, E.C. and Siperstein, G.N. (1970) Influence of contingent auditory stimulation upon non-nutritional suckle. In *Third Symposium on Oral Sensation and Perception: The Mouth of the Infant* (Bosma, J.F., ed.), pp. 313–334, Charles C. Thomas University Park Press
134. Eisenberg, R.B. (1976) *Auditory Competence in Early Life*, University Park Press
135. Sai, F.Z. (2005) The role of the mother's voice in developing mother's face preference: Evidence for intermodal perception at birth. *Infant Child Dev.* 14, 29–50
136. Walton, G.E. et al. (1992) Recognition of familiar faces by newborns. *Infant Behav. Dev.* 15, 265–269
137. Cupute, A.J., Palmer, F.B., Shupiro, B.K., Wuchtel, R.C., Ross, A. and Accurdo, P.J. (1984) Primitive reflex profile: a quantitation of primitive reflexes in infancy. *Dev. Med. Child Neurol.* 26, 375–383
138. Gesell, A. et al. (1934) *Infant Behavior: Its Genesis and Growth*, McGraw-Hill Book Company Inc.
139. Thalagala, N. et al. (2013) *Early Child Developmental Standards for Sri Lankan Infants and Toddlers*, UNICEF
140. Frankenburg, W.K. et al. (1992) The Denver II: A Major Revision and Restandardization of the Denver Developmental Screening Test. *Pediatrics* 89, 91–97
141. Gordon, M.B. (1929) The Moro embrace reflex in infancy: its incidence and significance. *Am. J. Dis. Child.* 38, 26–34
142. Touwen, B.C. (1971) A study on the development of some motor phenomena in infancy. *Dev. Med. Child Neurol.* 13, 435–446
143. Ramsay, J.O. and Silverman, B.W. (2006) *Functional Data Analysis*, Springer
144. Yao, F. et al. (2005) Functional data analysis for sparse longitudinal data. *J. Am. Stat. Assoc.* 100, 577–590
145. Jones, H. and Bayley, N. (1941) The Berkeley Growth Study. *Child Dev.* 12, 167–173
146. Leibold, L.J. and Werner, L.A. (2007) Infant Auditory Sensitivity to Pure Tones and Frequency-Modulated Tones. *Infancy* 12, 225–233
147. Fernald, A. et al. (1984) Expanded Intonation Contours in Mothers' Speech to Newborns. 20, 104–113
148. Telkemeyer, S. et al. (2009) Sensitivity of newborn auditory cortex to the temporal structure of sounds. *J. Neurosci.* 29, 14726–14733
149. Poeppel, D. (2003) The analysis of speech in different temporal integration windows: Cerebral lateralization as 'asymmetric sampling in time'. *Speech Commun.* 41 (150), 245–255
150. Pascalis, O. and Slater, A., eds (2003) *The Development of Face Processing in Infancy and Early Childhood: Current Perspectives*, Nova Science Publishers, (Hauppauge, NY, US